



Non bulbing type



A leafy, fast growing cross between Chinese cabbage and stubble turnip.

It exhibits high frost resistance and can be utilised within eight to ten weeks after sowing.

There is potential for regrowth.

Ratings

| | |
|-------------------------------------|---|
| Brassica Fodder Crop for CSS | <div style="width: 100%; height: 10px; background-color: green;"></div> |
| Dry Matter Yield | <div style="width: 60%; height: 10px; background-color: green;"></div> |
| Crude Protein | <div style="width: 80%; height: 10px; background-color: green;"></div> |
| Digestibility | <div style="width: 70%; height: 10px; background-color: green;"></div> |
| Metabolisable Energy | <div style="width: 70%; height: 10px; background-color: green;"></div> |
| Bulb Size | <div style="width: 20%; height: 10px; background-color: green;"></div> |
| Bolting Resistance | <div style="width: 60%; height: 10px; background-color: green;"></div> |
| Winter Hardiness | <div style="width: 70%; height: 10px; background-color: green;"></div> |

| Seed Rate | Pack Size | S |
|--------------|-----------|----|
| 5 - 7.5kg/ha | 5kg | TI |

Stock should be introduced gradually over a two week period and an area of grassland should be available for animals to return to; water and hay or straw should also be made available. Please contact your supplier for further guidance.

| Stubble Turnips Yield and Feed Quality | |
|--|--------------------------|
| Average dry matter yield | 3.5 - 4.5 tonnes/ha |
| Average fresh yields | 38 – 45 tonnes/ha |
| Crude protein | 17 - 18% (mainly leaves) |
| Digestibility value | 68 - 70 D |
| Dry matter | 8 - 9% |
| Metabolisable energy | 11 MJ/kg DM |
| Sugars in DM | 55% |